

BENIL^{Tablets}

Glibenclamide B.P.

بنیل گولیاں
گلیبن کلیمین کا ایسی ڈی۔ پی۔

Composition:

Each tablets contains:

Glibenclamide B.P.5mg

BENIL is the name given to the formula 1-[[4-[2-[(5-chloro - 2 - methoxybenzoyl) amino]ethyl]phenyl]sulphonyl] 3-cyclohexyiurea, chemically known as Glibenclamide.

Indications:

BENIL is indicated for Diabetes Mellitus. It is an orally active hypoglycaemic agent which can reduce the blood-sugar level, reduce or abolish glycosurea and improve metabolic control.

Contra-Indications:

BENIL is contra indicated in impaired renal or hepatic function, andrendortical insufficiency and during pregnancy.

Precautions:

The hypoglycaemic effects of benil may be enhanced by alcohol, monoamine oxidase inhibitors, beta-adrenergic blocking agents and some sulphonamides. The hypoglycaemic effects may be diminished by adrenaline, corticosteroids and thiazide diuretics. If any symptoms of hypoglycaemia appears during the treatment with the physician should be informed immediately. The physician should also be informed if any other drug is being used simultaneously.

Dosage:

2.5 to 10 mg once daily. Daily dosage of upto 2 tablets may be taken after breakfast. When the daily dosage exceeds 2 tablets, the excess is to be taken with the evening meals. The tablets are scored to be easily halved. The dosage and time of intake of **Benil** must be followed as directed by the physician. The change from another antidiabetic treatment to **Benil** should always be carried out under the guidance of a physician. A person changing physician should explain his diabetic condition to the new physician.

Presentation:

BENIL 5mg tablets are supplied in a carton (3x20) Blister Pack.



Manufactured by:
REKO PHARMACAL (Pvt.) LTD.
13-km, Multan Road, Lahore-Pakistan.
www.rekopharmacal.com

